

Pain and Depression: A Neurobiological Perspective of Their Relationship

Changsu Han¹ and Chi-Un Pae^{2,3} ✉

¹Department of Psychiatry, Korea University College of Medicine, Seoul, Republic of Korea

²Department of Psychiatry, Bucheon St. Mary's Hospital, The Catholic University of Korea College of Medicine, Bucheon, Republic of Korea

³Department of Psychiatry and Behavioral Sciences, Duke University Medical Center, Durham, NC, USA

Remarkable progresses have been achieved regarding the understanding of the neurobiological bases of pain and depression. The principal role of neurotransmitters, neuromodulators, and neurohormones has been proposed in the development of pain and depression. With the progression of molecular biology, an intricate interaction among biological factors accountable to the development and management of pain and depression has been also shown in a numerous preclinical and clinical researches. This mini-review will briefly describe the current issues and future research direction for better understanding of the relationship between pain and depression.

Psychiatry Investig 2015;12(1):1-8

Key Words Depression, Pain, Relationship, Pathophysiology, Symptomatology, Treatment.

INTRODUCTION

Depression and pain commonly occur together and both can have detrimental effects on patient health, personal and social function, and overall quality of life. Pain is a controversial syndrome and its clear phenomenology, aetiology, pathogenesis and treatment remain a matter of controversy.¹ Numerous studies indicate that patients with pain are at increased risk for a lifetime history of depression, as compared to the normal population.²

According to the World Health Organisation study in primary care settings across the world, approximately one-fifth of all primary care patients suffer from persistent debilitating pain, and they are four times more likely to have co-morbid anxiety or depression than pain-free primary care patients.³

Similarly, depressed patients are also associated with more pain complaints and greater impairment in functional capacity. In addition, the risk of depression is greater when the pain is

more diffuse, as indicated by the number of painful sites, and has a greater effect on the quality of life.^{4,5}

In addition, serotonin and norepinephrine were principally investigated since these neurotransmitters are located in the main pain pathway as well as preclinical and clinical evidence have suggested a significant role of such neurotransmitters regarding the development and treatment of pain.⁶ In fact, a number of psychotropic agents such as selective serotonin reuptake inhibitors (SSRIs), noradrenergic and specific serotonin receptor antagonist (NaSSa), serotonin-norepinephrine reuptake inhibitors (SNRIs) have been actively tried for the treatment of pain disorders or painful somatic symptoms regardless of patients' diagnosis, all of which are involved in serotonin and norepinephrine neurotransmission.⁶

This brief perspective will provide a summary of intricate relationship between pain and depression as a further understanding on the role of pain in relation with the pathophysiology, clinical manifestation, and proper management of depression.

COMMON UNDERSTANDING ON ETIO-PATHOLOGIC MECHANISM

Overall findings

The precise etio-pathological mechanism of pain remains still unclear. A number of clinical and biological factors have

Received: December 30, 2013 **Revised:** February 13, 2014

Accepted: March 13, 2014 **Available online:** January 12, 2015

✉ **Correspondence:** Chi-Un Pae, MD, PhD

Department of Psychiatry, Bucheon St. Mary's Hospital, The Catholic University of Korea College of Medicine, 327 Sosa-ro, Wonmi-gu, Bucheon 420-717, Republic of Korea

Tel: +82-32-430-2140, **Fax:** +82-32-340-2255, **E-mail:** pae@catholic.ac.kr

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

been demonstrated to be involved in the development of pain. For instance, altered neuromodulation,⁷ abnormalities of the immune-genetic system (i.e., alterations in pro-inflammatory cytokines⁸ and their gene expression),⁹ alteration in melatonin,¹⁰ aberration in ³H-imipramine binding sites on platelets,¹¹ the oxidative stress reaction,⁹ a disturbance of neurotransmitters,¹² stress susceptibility,¹³ alterations of hypothalamus-pituitary-adrenal axis function,¹⁴ and cognitive vulnerability¹⁵ are known to at least partly play a role in the manifestation of pain. In addition, a number of psychosocial factors (i.e., self esteem, self-efficacy, anger, sexual abuse, personality profile, early childhood abuse) are also crucially involved in the development of pain.¹⁶⁻¹⁸ Intriguingly all of which clinical and biological factors aforementioned are also common in the development of depression.

Genetic factors

Patients with chronic pain have more first degree relatives with depression compared with the general population.^{3,19} Significantly higher rates of depression have been found in family members even for patients with chronic pain without a personal history of depression.^{3,11} According to a previous human genetic study,²⁰ the galanin-2 and mu opioid receptors are plausible candidates for mediating the effects of pain on mood, as they show graphic patterns consistent with a pain-gene interaction, although such specific effects cannot be conclusive due to the modest sample size of the study. In addition, a recent study found that serotonin 1A receptor (5HTR1A) and serotonin 2A receptor gene promoter variations have gender-dependent modulatory effects on depression and physical function in patients with pain. Furthermore, that study demonstrated that pain after lumbar surgery modulates the association between 5HT gene polymorphisms and depression.²¹ These findings suggest that pain sensitivity, the stress response and mood regulation share common familial-genetic factors and support the hypothesis that depression and pain are genetically related.

Pain pathways and biochemical findings

The CNS pathway responsible for inhibition of pain sensation includes projections from brainstem nuclei to the spinal cord dorsal horn via the dorsolateral funiculus (DLF). More specifically, DLF fibers are comprised of serotonergic projections from the raphe nuclei, dopaminergic projections from the ventral tegmental area (VTA), and noradrenergic projections from the locus coeruleus.^{22,23} These descending fibers suppress pain transmission at the nociceptive spinal cord neurons presumably by hyperpolarizing afferent sensory neurons using endogenous opioids, or serotonin and norepinephrine as principal inhibitory mediators.^{22,23}

Activity induced in the nociceptors and nociceptive pain pathways is not pain, but is nociception, and pain perception is formed in the central nervous system as a very interactive phenomenon through central and peripheral systems. The pathophysiology of acute pain is fairly clear, but chronic pain is still a great enigma, and includes a number of neurotransmitters (GABA, glutamate, norepinephrine, serotonin, neurokinin 1, nitric oxide, substance P, glycine and opioids).²⁴ It has been proposed that central and peripheral pain mechanisms may be altered in patients with pain, resulting in amplified pain signals.²⁵ Current theories assume that continuous stimulation of the pain pathways (C-fibres) leads to over-activation of N-methyl-D-aspartate receptors in the dorsal horn of the spinal cord.²⁶ In the first human study investigating that repetitive stimulation enhances a bidirectional mechanism of secondary hyperalgesia due to viscerosomatic facilitation in patients with irritable bowel syndrome (IBS),²⁷ the IBS patients were tested by repetitive stimulation of rectal and somatic tissues to induce central sensitization as well as an NMDA receptor antagonist (dextromethorphan) versus placebo. According to the results, enhanced pain sensitivity was nearly completely prevented by administration of dextromethorphan, suggesting that pharmacological manipulation of NMDA receptor mechanisms is essentially involved in modulation of central sensitization for processing of pain.

Increased secretion of excitatory neuromodulators, such as substance P, and psychological factors further escalate such pain perception. The resulting state is referred to as "central sensitisation" wherein intense pain is produced from innocuous stimuli.²⁸ For example, patients with pain show reduced blood flow in the caudate and thalamic areas involved in the central processing of noxious signals²⁹ and elevated concentrations of cerebrospinal fluid substance P³⁰ and brain-derived neurotrophic factor.³¹ In a recent study,³² enhanced central pain sensitivity at both ends of the spinal neuroaxis was found in patients with fibromyalgia and therefore widespread central sensitisation. These findings are commonly observed in patients with depression as well.^{33,34} Table 1 represents the potential similarities in biological markers between pain disorders and depression.

The principal biochemical basis for pain and depression has focused on the neurotransmitters serotonin and norepinephrine. Both the serotonin and norepinephrine pathways in the brain and their associated symptoms have been determined. As described before, both pathways originate in the brain stem nuclei and project to multiple brain regions such as somatosensory cortex, intralaminar nuclei and ventral posterior nuclei of the thalamus.³⁵ In the context of pain regulation, peripheral pathways descend projections from the brain stem to the spinal cord. The serotonin and norepinephrine descending

Table 1. Potential similarities in biological markers between pain disorders and depression^{74,75)}

	Pain disorders	Depression
Neurotransmitters and those-related markers		
Tryptophan in CSF and plasma	Low	Low
5-HT in serum	Low	Low with inconsistency
5-HT in platelet	Low	Low
5-HIAA	Low	Low
MHPG	Low	Low
HVA	Low	Low
Hormones		
Growth hormone	Low	Low
Estrogen	Low	Low
Tri-iodothyroxine	Low	Low
Blunted TSH response to TRH	+	+
Melatonin	Low	Low
Neuroendocrinologic markers		
Substance P in CSF and plasma	Increase	Increase
Non-suppression of dexamethasone	±	+
Plasma endorphin	Low	Low
Corticotropin-releasing factor	++	++
Glucocorticoids	++	++
Neurotrophic markers		
Total antioxidant capacity	Low	Low
Brain-derived neurotrophic factor	Low	Low
Nerve growth factor	Low	Low
Erythrocyte levels of n-3 PUFA	Low	Low
Immune-modulators		
IL-1β	Increase	Increase
IL-2	Increase	Increase
IL-6	Increase	Increase
IL-8	Increase	Increase
IL-10	Increase	Increase
C-reactive protein	Increase	Increase
Tumor necrosis factor-alpha	Increase	Increase
CD40 ligand	Increase	Increase
Antidepressant effects	++	+++

neurons project from the brain stem into the spinal cord's dorsal horn, where a set of biochemical cascades takes place involving other types of neurotransmitters for controlling pain.³⁵ The Figure 1 represents schematic illustration of the pain pathway.

Disturbed cognition

Many cognitive variables have been shown to mediate the pain-depression association, such as pessimism, perceived lo-

cus of control, fear avoidance beliefs, self-efficacy, and perceived social support.^{36,37} Perhaps the best studied and potentially most relevant to the pain-depression relationship is catastrophizing that is literally to view or present a situation as considerably worse than it actually is. Specifically, in psychological researches, it refers to cognitive and emotional processes encompassing magnification of pain-related stimuli, feelings of helplessness, and a generally pessimistic orientation. The term was initially introduced by Albert Ellis and subse-




Figure 1. The physiological pathway of pain. FC: frontal cortex, PSC: primary sensory cortex, HT: hypothalamus, CC: cingulate cortex, AMD: amygdala, MTN: medial thalamic nuclei, LTN: lateral thalamic nuclei, FNE: free nerve ending. The whole process of perception of pain is following: Transduction → Transmission (central sensitization) → Modulation → Perception.

quently adapted by Aaron Beck to describe a mal-adaptive cognitive style employed by patients with anxiety and depressive disorders.³⁸ Hence, the catastrophizing is currently considered a cognitive error where one assumes the worst possible outcome will occur.³⁹ It has been associated with pain and depression separately, and they co-occur as well.³⁷ Depression and catastrophizing has been found to be associated with higher pain severity and functional incapacity, conversely, researches also propose that those who catastrophize their pain tend to feel helpless, dreadful, and even worthless.³⁷ Recent research supporting such viewpoint studied 669 older adults with chronic pain and found that catastrophization mediated the relationship between pain intensity and depression.⁴⁰

Overlapping clinical symptoms

Patients with pain and depression have important similarities, as reflected in elevated levels of depressed mood,⁴¹ alexithymia,⁴² anger,⁴³ decreased energy and fatigue,⁴⁴ cognitive defect,⁴⁵ increased vulnerability to stress,⁴⁶ heart rate variability⁴⁷ and sleep disturbance (decreased REM latency).⁴⁸ Furthermore, the severity of pain is highly associated with some depression-related symptoms and impairments such as depressed mood, fatigue, functional disability, suicidal ideation, poor daily health status and more somatic symptoms.^{45,49,50}

Effects of antidepressants and cognitive therapy

Despite no evidence for the superiority of one class of antidepressants in the treatment of pain, contemporary antidepressants involving serotonin and norepinephrine activities such as venlafaxine, duloxetine and milnacipran have consistently shown the promising effect for controlling pain among vari-

able psychotropic agents.

However, recently, in a number of clinical trials with dopaminergic agents have also been conducted and their beneficial effects for controlling pain⁵¹ and depression¹⁹ have been demonstrated. A recent study investigated whether or not pain-related behavioural depression is mediated by activation of endogenous κ -opioid systems and subsequent depression of mesolimbic dopamine release.⁵² That study found a crucial role of dopamine neurotransmission in terms of pain-related depression of behaviour, pain-related depression of mesolimbic dopamine release and role of endogenous dynorphin/ κ -opioid receptor systems: 1) the acid noxious stimulus also depressed extracellular levels of the neurotransmitter dopamine in nucleus accumbens; supporting the fact that depression of mesolimbic dopamine release may contribute to negative affective dimensions of pain; 2) acid-induced depression of dopamine release was blocked by both NSAID and opioid analgesics, indicating the potential relationship of opioid with affective dimensions of pain, and 3) alteration of mesocorticolimbic prodynorphin expression by noxious stimulus.⁵² In addition, intriguingly, milnacipran does not significantly modulate the activity of segmental spinal nociceptive neurons or influence the diffuse noxious inhibitory control activity or the documented thermoalgebraic allodynia, but the supraspinal analgesic effect might be involved in the mechanism of milnacipran for controlling pain.⁵³ For instance, dopaminergic agents for pain conditions like fibromyalgia and neuropathic pain, showed varied results. In a study with pramipexole, a dopamine receptor agonist, for the treatment of fibromyalgia, the primary outcome was improvement in the pain score [10-cm visual analog scale (VAS)] at 14 weeks.⁵¹ Compared with the

placebo group, patients-treated with pramipexole experienced gradual and more significant improvement in measures of pain. At the endpoint, the VAS pain score decreased 36% in the pramipexole arm and 9% in the placebo arm, indicating a huge magnitude of difference of 27%. Forty-two percent of patients receiving pramipexole and 14% of those receiving placebo achieved $\geq 50\%$ decrease in pain. This study has clearly established the efficacy and safety of pramipexole in a subset of patients with fibromyalgia.⁵¹ Bupropion, a representative antidepressant acting on dopamine modulation, has been also tested in the management of pain.⁵⁴ In an outpatient-based (n=42), randomized, double-blind, placebo-controlled, crossover study consisted of two phases (12 weeks), bupropion significantly reduced the mean average pain score by 1.7 points to 4.0 ($p < 0.001$), while placebo treatment failed to do so. Such effect started to be significant at week 2 and remained throughout weeks 3 through 6 ($p < 0.001$). A significant decrease in interference of pain on quality of life was also observed, while it was not shown in placebo treatment.

In this context, it appears reasonable that all three major neurotransmitters are related to the pathophysiology of depression/pain and, as such, are relevant targets for pharmacological intervention. Triple reuptake inhibitors under development, which enhance neurotransmission of all three systems, should be a potential next-generation therapeutic agent to provide more reliable efficacy and a quicker therapeutic effect.⁵⁵ Hence, enhancing the current understanding of the exact action mechanism of antidepressants on pain pathways should also be simultaneously examined to expand our knowledge, thus leading to better pharmacological management and clinical outcomes for such patients.

Psychological interventions took a main position in modern pain management practice and have also been a recommended option of a modern pain treatment service. A number of systematic reviews and meta-analyses have been conducted and provided the evidence of psychological interventions for the management of pain in clinical practice.^{56,57} Among such psychological interventions, cognitive behavioural therapy (CBT) with a focus on cognitive coping strategies and behavioural rehearsal has demonstrated the strongest evidence. However, currently available most evidence is for treatments of adult pain but few for adolescent and older patients with chronic pain.⁵⁸ According to such well-researched systematic reviews, beneficial effects without any harmful effects by using any medications were prominently evident with CBT.^{56,59} It has been proven that substantial improvement in pain, mood, and functional incapacity which were not able to be achieved by any other form of treatment, was also remarkable. However, the overall effect sizes of CBT for adults, across all trials, were modest.⁵⁶

DISCUSSION

Despite the fact that pain and depression potentiate each other clinically, and an increasing number of neuroimaging studies show that physical pain and depression involve the insular cortex,⁶⁰ a recent functional imaging study⁶¹ clearly suggested that they may involve different cortical pain areas. According to that study, patients with depression show less activation of the operculum and the secondary somatosensory cortex, as compared to that in patients with pain, whereas patients with pain demonstrate increased activation in the




Figure 2. The association of pain and depression in pathophysiology and symptomatology.

anterior insula and parietal operculum. In addition, in all participants sensitised to pain, there was an association with higher activation levels in the thalamus, amygdala, mid-cingulate cortex and sensory and motor areas; however, depressed patients showed significantly less activation in midbrain and brainstem areas. Hence, newer imaging techniques, possibly proving the existence of different neuronal correlates noninvasively between pain and depression in terms of brain metabolism and hemodynamics during acute and chronic phases, would also be helpful in understanding both conditions.

Emerging evidence suggests that dopamine may be another crucial factor associated with symptoms of pain and depression, whereas previous pain and depression studies have focused mainly on serotonin and norepinephrine. In fact, some anecdotal reports have proposed that the VTA is related to the processing of nociceptive information as well as control of motor function. In fact, in a recent rodent study,⁶² VTA lesions enhanced the occurrence of self-injury behaviour, while activation of the VTA by electrical stimulation after a nociceptive stimulus, but not before, facilitated the analgesic process, indicating that the VTA should play a key role in the processing and modulation of persistent pain information. Fian and Smith⁶³ have also proposed that the homeostatic regulation of tonic and phasic dopamine may contribute to both pain and depression. Hence, attention to the role of dopamine in pain and depression would help clinicians understand both clinical conditions more fully.

Although increasing evidence indicates that the relationship between both clinical conditions might be bidirectional,⁶⁴ the question remains whether the specific neurobiological changes cause either pain or depression or are merely a consequence of each condition. Interestingly, some researchers proposed the concept of chronic pain as a psychological disorder since there is some support to view chronic pain as the prime expression of a muted depressive state;⁶⁵ however, it has been remained a vague entity due to a lack of continuous supporting data for this notion. Integrative and comprehensive approaches for revealing the association between pain and depression would be more straightforward and reasonable based on currently available findings.

Depression and pain share clinical manifestations, neurobiological pathways, neurotransmitters, hormones, stressors, neurotrophic factors and pro-inflammatory cytokines, which have implications for the development and treatment of both simultaneously. The final common pathway in which chronic pain and depression may be ultimately associated should be when both arise out of a common underlying process.

Another huge interesting point is that we need to consider the relationship between suicide and pain since the suicide rate is higher in patients with depression as well as with pain, in-

dicating that pain could be a potentially common mediator or risk factor for both clinical conditions. In fact, clinical data consistently suggest that patients with medico-surgical conditions including malignancies and cardiovascular diseases causing serious disability of patients or simple but chronic headaches are at an increased risk of committing suicide.⁶⁶⁻⁶⁹ For instance, according to large-scale studies, patients with migraine are from 2.2 to 4.0 times more likely to have depression⁷⁰ and as well as more likely to have suicide mortality (odds ratio=1.68).⁶⁹ In addition, patients with migraine have been found to be associated with more increased risk of developing suicidal ideation and committing suicide attempts when they are comorbid with depression.⁷⁰ Hence, it should be also intriguing for clinicians to investigate how pain and depression are interactive and implicated in the development and prevention of suicide.

Next researches should also try to clearly establish the relationship between acute and chronic pain with depression. In fact, acute and chronic pain are different clinical entities. Acute pain is usually provoked by a specific disease or injury and is self-limited, while chronic pain may be considered as a disease state and also arise from psychological states.^{71,72} Chronic pain has been proposed to be the consequence of persistent impulses in somatosensory pathways with related CNS changes (i.e., altered local brain chemistry and functional reorganization due to central plasticity).⁷³ In addition, the management and treatment of acute and chronic pain patients are also substantially different due to its basic differences in clinical course and symptoms. However, there has been a paucity of data concerning a differential relationship between acute and chronic pain and depression till today. Hence this clinical issue warrants well-designed future studies.

Contemporary findings suggest that understanding the brain is a critical and indispensable component of pain and depression research involved in explaining the ways in which biological, psychological, behavioural and social factors influence both pain and depression, and offers a way to the optimal therapeutic approach (Figure 2).²⁴ Psychosocial and behavioural factors play a significant role in the experience, maintenance, and exacerbation of pain.³⁷ CBT alone or within the context of an interdisciplinary pain rehabilitation program has the greatest empirical evidence for success.³⁷ As none of the most commonly prescribed treatment regimens are sufficient to eliminate pain, a more realistic approach will likely combine pharmacological, physical, and psychological components tailored to each patient's needs.⁵⁹

Therefore, a bio-psycho-social model that integrates aetiology, evaluation and management of depression and pain concurrently is mandatory for better care of such patients. We must promote further theoretical, methodological and tech-

nical advances to clarify the precise relationship between these conditions, which may also lead to more effective biological and psychological therapies for such patients.

Acknowledgments

This study was partially funded by the Ministry of Health and Welfare, Korea (HI12C0003).

REFERENCES

1. Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, et al. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis Rheum* 1990;33:160-172.
2. Ratcliffe GE, Enns MW, Belik SL, Sareen J. Chronic pain conditions and suicidal ideation and suicide attempts: an epidemiologic perspective. *Clin J Pain* 2008;24:204-210.
3. Lepine JP, Briley M. The epidemiology of pain in depression. *Hum Psychopharmacol* 2004;19(Suppl 1):S3-S7.
4. Pae CU, Luyten P, Marks DM, Han C, Park SH, Patkar AA, et al. The relationship between fibromyalgia and major depressive disorder: a comprehensive review. *Curr Med Res Opin* 2008;24:2359-2371.
5. Pae CU, Marks DM, Patkar AA, Masand PS, Luyten P, Serretti A. Pharmacological treatment of chronic fatigue syndrome: focusing on the role of antidepressants. *Expert Opin Pharmacother* 2009;10:1561-1570.
6. Sansone RA, Sansone LA. Pain, pain, go away: antidepressants and pain management. *Psychiatry (Edgmont)* 2008;5:16-19.
7. Leone M. Deep brain stimulation in headache. *Lancet Neurol* 2006;5:873-877.
8. Bai YM, Chiou WF, Su TP, Li CT, Chen MH. Pro-inflammatory cytokine associated with somatic and pain symptoms in depression. *J Affect Disord* 2014;155:28-34.
9. Burke NN, Geoghegan E, Kerr DM, Moriarty O, Finn DP, Roche M. Altered neuropathic pain behaviour in a rat model of depression is associated with changes in inflammatory gene expression in the amygdala. *Genes Brain Behav* 2013;12:705-713.
10. Blumer D, Zorick F, Heilbronn M, Roth T. Biological markers for depression in chronic pain. *J Nerv Ment Dis* 1982;170:425-428.
11. Magni G, Andreoli F, Arduino C, Arsie D, Ceccherelli F, Ambrosio F, et al. Modifications of [3H]imipramine binding sites in platelets of chronic pain patients treated with mianserin. *Pain* 1987;30:311-320.
12. Nicolson SE, Caplan JP, Williams DE, Stern TA. Comorbid pain, depression, and anxiety: multifaceted pathology allows for multifaceted treatment. *Harv Rev Psychiatry* 2009;17:407-420.
13. Sachs-Ericsson N, Cromer K, Hernandez A, Kendall-Tackett K. A review of childhood abuse, health, and pain-related problems: the role of psychiatric disorders and current life stress. *J Trauma Dissociation* 2009;10:170-188.
14. Muhtz C, Rodriguez-Raecke R, Hinkelmann K, Moeller-Bertram T, Kiefer F, Wiedemann K, et al. Cortisol response to experimental pain in patients with chronic low back pain and patients with major depression. *Pain Med* 2013;14:498-503.
15. Lee EJ, Wu MY, Lee GK, Cheing G, Chan F. Catastrophizing as a cognitive vulnerability factor related to depression in workers' compensation patients with chronic musculoskeletal pain. *J Clin Psychol Med Settings* 2008;15:182-192.
16. Scott KM, Von Korff M, Angermeyer MC, Benjet C, Bruffaerts R, de Girolamo G, et al. Association of childhood adversities and early-onset mental disorders with adult-onset chronic physical conditions. *Arch Gen Psychiatry* 2011;68:838-844.
17. Jang JM, Park JI, Oh KY, Lee KH, Kim MS, Yoon MS, et al. Predictors of suicidal ideation in a community sample: roles of anger, self-esteem, and depression. *Psychiatry Res* 2014;216:74-81.
18. Friberg O, Martinsen EW, Martinussen M, Kaiser S, Overgard KT, Rosenvinge JH. Comorbidity of personality disorders in mood disorders: a meta-analytic review of 122 studies from 1988 to 2010. *J Affect Disord* 2014;152-154:1-11.
19. Corrigan MH, Denahan AQ, Wright CE, Ragual RJ, Evans DL. Comparison of pramipexole, fluoxetine, and placebo in patients with major depression. *Depress Anxiety* 2000;11:58-65.
20. Max MB, Wu T, Atlas SJ, Edwards RR, Haythornthwaite JA, Bollettino AF, et al. A clinical genetic method to identify mechanisms by which pain causes depression and anxiety. *Mol Pain* 2006;2:14.
21. Lebe M, Hasenbring MI, Schmieder K, Jetschke K, Harders A, Epplen JT, et al. Association of serotonin-1A and -2A receptor promoter polymorphisms with depressive symptoms, functional recovery, and pain in patients 6 months after lumbar disc surgery. *Pain* 2013;154:377-384.
22. Dafny N. Pain Modulation and Mechanisms In Section 2: Sensory Systems. In: Byrne JH, Editor. *Neuroscience Online*. Houston: The University of Texas Health Science Center at Houston; 2008.
23. Marks DM, Patkar AA, Masand PS, Pae CU. Does pregabalin have neuropsychotropic effects?: a short perspective. *Psychiatry Investig* 2009;6:55-58.
24. Bras M, Dordevic V, Gregurek R, Bulajic M. Neurobiological and clinical relationship between psychiatric disorders and chronic pain. *Psychiatr Danub* 2010;22:221-226.
25. Bendtsen L, Norregaard J, Jensen R, Olesen J. Evidence of qualitatively altered nociception in patients with fibromyalgia. *Arthritis Rheum* 1997;40:98-102.
26. Bennett M. Emerging concepts in the neurobiology of chronic pain: evidence of abnormal sensory processing in fibromyalgia. *Mayo Clin Proc* 1999;74:385-398.
27. Verne GN, Price DD, Callam CS, Zhang B, Peck J, Zhou Q. Viscerosomatic facilitation in a subset of IBS patients, an effect mediated by N-methyl-D-aspartate receptors. *J Pain* 2012;13:901-909.
28. Doubell TP, Mannion RJ, Woolf CJ. The Dorsal Horn: State-Dependent Sensory Processing, Plasticity and the Generation of Pain. In: Wall PD, Melzack R, Editors. *Textbook of pain*. Edinburgh: Churchill Livingstone, 1999, p.165-181.
29. Mountz JM, Bradley LA, Modell JG, Alexander RW, Triana-Alexander M, Aaron LA, et al. Fibromyalgia in women. Abnormalities of regional cerebral blood flow in the thalamus and the caudate nucleus are associated with low pain threshold levels. *Arthritis Rheum* 1995;38:926-938.
30. Russell IJ, Orr MD, Littman B, Vipraio GA, Alboukrek D, Michalek JE, et al. Elevated cerebrospinal fluid levels of substance P in patients with the fibromyalgia syndrome. *Arthritis Rheum* 1994;37:1593-1601.
31. Laske C, Stransky E, Eschweiler GW, Klein R, Wittorf A, Leyhe T, et al. Increased BDNF serum concentration in fibromyalgia with or without depression or antidepressants. *J Psychiatr Res* 2007;41:600-605.
32. Staud R, Robinson ME, Price DD. Temporal summation of second pain and its maintenance are useful for characterizing widespread central sensitization of fibromyalgia patients. *J Pain* 2007;8:893-901.
33. Blier P, Gobbi G, Haddjeri N, Santarelli L, Mathew G, Hen R. Impact of substance P receptor antagonism on the serotonin and norepinephrine systems: relevance to the antidepressant/anxiolytic response. *J Psychiatry Neurosci* 2004;29:208-218.
34. Zelman DC, Howland EW, Nichols SN, Cleeland CS. The effects of induced mood on laboratory pain. *Pain* 1991;46:105-111.
35. Jann MW, Slade JH. Antidepressant agents for the treatment of chronic pain and depression. *Pharmacotherapy* 2007;27:1571-1587.
36. Philips HC. Imagery and likelihood cognitive bias in pain. *Behav Cogn Psychother* 2013;1-15.
37. Goesling J, Clauw DJ, Hassett AL. Pain and depression: an integrative review of neurobiological and psychological factors. *Curr Psychiatry Rep* 2013;15:421.
38. Quartana PJ, Campbell CM, Edwards RR. Pain catastrophizing: a critical review. *Expert Rev Neurother* 2009;9:745-758.

39. Hassett AL, Cone JD, Patella SJ, Sigal LH. The role of catastrophizing in the pain and depression of women with fibromyalgia syndrome. *Arthritis Rheum* 2000;43:2493-2500.
40. Nicholas MK, Asghari A, Blyth FM, Wood BM, Murray R, McCabe R, et al. Self-management intervention for chronic pain in older adults: a randomised controlled trial. *Pain* 2013;154:824-835.
41. Burckhardt CS, O'Reilly CA, Wiens AN, Clark SR, Campbell SM, Bennett RM. Assessing depression in fibromyalgia patients. *Arthritis Care Res* 1994;7:35-39.
42. Saariaho AS, Saariaho TH, Mattila AK, Karukivi MR, Joukamaa MI. Alexithymia and depression in a chronic pain patient sample. *Gen Hosp Psychiatry* 2013;35:239-245.
43. Sayar K, Gulec H, Topbas M. Alexithymia and anger in patients with fibromyalgia. *Clin Rheumatol* 2004;23:441-448.
44. Moldovan I, Cooray D, Carr F, Katsaros E, Torralba K, Shinada S, et al. Pain and depression predict self-reported fatigue/energy in lupus. *Lupus* 2013;22:684-689.
45. Zakoscielna KM, Parmelee PA. Pain variability and its predictors in older adults: depression, cognition, functional status, health, and pain. *J Aging Health* 2013;25:1329-1339.
46. Martinez-Lavin M. Biology and therapy of fibromyalgia. Stress, the stress response system, and fibromyalgia. *Arthritis Res Ther* 2007;9:216.
47. Toro-Velasco C, Arroyo-Morales M, Fernandez-de-Las-Penas C, Cleland JA, Barrero-Hernandez FJ. Short-term effects of manual therapy on heart rate variability, mood state, and pressure pain sensitivity in patients with chronic tension-type headache: a pilot study. *J Manipulative Physiol Ther* 2009;32:527-535.
48. Wilson KG, Eriksson MY, D'Eon JL, Mikail SF, Emery PC. Major depression and insomnia in chronic pain. *Clin J Pain* 2002;18:77-83.
49. Olie E, Guillaume S, Jaussent I, Courtet P, Jollant F. Higher psychological pain during a major depressive episode may be a factor of vulnerability to suicidal ideation and act. *J Affect Disord* 2010;120:226-230.
50. Jeon HJ, Lee JY, Lee YM, Hong JB, Won SH, Cho SJ, et al. Unplanned versus planned suicide attempters, precipitants, methods, and an association with mental disorders in a Korea-based community sample. *J Affect Disord* 2010;127:274-280.
51. Holman AJ, Myers RR. A randomized, double-blind, placebo-controlled trial of pramipexole, a dopamine agonist, in patients with fibromyalgia receiving concomitant medications. *Arthritis Rheum* 2005;52:2495-2505.
52. Leitl MD, Onvani S, Bowers MS, Cheng K, Rice KC, Carlezon WA Jr, et al. Pain-related depression of the mesolimbic dopamine system in rats: expression, blockade by analgesics, and role of endogenous k-opioids. *Neuropsychopharmacology* 2014;39:614-624.
53. Matthey A, Cedraschi C, Piguet V, Besson M, Chabert J, Daali Y, et al. Dual reuptake inhibitor milnacipran and spinal pain pathways in fibromyalgia patients: a randomized, double-blind, placebo-controlled trial. *Pain Physician* 2013;16:E553-E562.
54. Semenchuk MR, Sherman S, Davis B. Double-blind, randomized trial of bupropion SR for the treatment of neuropathic pain. *Neurology* 2001;57:1583-1588.
55. Marks DM, Pae CU, Patkar AA. Triple reuptake inhibitors: a premise and promise. *Psychiatry Investig* 2008;5:142-147.
56. Eccleston C, Morley SJ, Williams AC. Psychological approaches to chronic pain management: evidence and challenges. *Br J Anaesth* 2013;111:59-63.
57. Williams AC, Eccleston C, Morley S. Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane Database Syst Rev* 2012;11:CD007407.
58. Eccleston C, Palermo TM, Fisher E, Law E. Psychological interventions for parents of children and adolescents with chronic illness. *Cochrane Database Syst Rev* 2012;8:CD009660.
59. Turk DC, Swanson KS, Tunks ER. Psychological approaches in the treatment of chronic pain patients--when pills, scalpels, and needles are not enough. *Can J Psychiatry* 2008;53:213-223.
60. Mutschler I, Ball T, Wankerl J, Strigo IA. Pain and emotion in the insular cortex: evidence for functional reorganization in major depression. *Neurosci Lett* 2012;520:204-209.
61. Rodriguez-Raecke R, Ihle K, Ritter C, Muhtz C, Otte C, May A. Neuronal differences between chronic low back pain and depression regarding long-term habituation to pain. *Eur J Pain* 2014;18:701-711.
62. Sotres-Bayon F, Torres-Lopez E, Lopez-Avila A, del Angel R, Pellicer F. Lesion and electrical stimulation of the ventral tegmental area modify persistent nociceptive behavior in the rat. *Brain Res* 2001;898:342-349.
63. Finan PH, Smith MT. The comorbidity of insomnia, chronic pain, and depression: dopamine as a putative mechanism. *Sleep Med Rev* 2013;17:173-183.
64. Raphael KG, Janal MN, Nayak S, Schwartz JE, Gallagher RM. Familial aggregation of depression in fibromyalgia: a community-based test of alternate hypotheses. *Pain* 2004;110:449-460.
65. Blumer D, Heilbronn M. Chronic pain as a variant of depressive disease: the pain-prone disorder. *J Nerv Ment Dis* 1982;170:381-406.
66. Beghi M, Rosenbaum JF, Cerri C, Cornaggia CM. Risk factors for fatal and nonfatal repetition of suicide attempts: a literature review. *Neuropsychiatr Dis Treat* 2013;9:1725-1736.
67. Robinson D, Renshaw C, Okello C, Moller H, Davies EA. Suicide in cancer patients in South East England from 1996 to 2005: a population-based study. *Br J Cancer* 2009;101:198-201.
68. Fall K, Fang F, Mucci LA, Ye W, Andren O, Johansson JE, et al. Immediate risk for cardiovascular events and suicide following a prostate cancer diagnosis: prospective cohort study. *PLoS Med* 2009;6:e1000197.
69. Ilgen MA, Kleinberg F, Ignacio RV, Bohnert AS, Valenstein M, McCarthy JF, et al. Noncancer pain conditions and risk of suicide. *JAMA Psychiatry* 2013;70:692-697.
70. Pompili M, Di Cosimo D, Innamorati M, Lester D, Tatarelli R, Martelletti P. Psychiatric comorbidity in patients with chronic daily headache and migraine: a selective overview including personality traits and suicide risk. *J Headache Pain* 2009;10:283-290.
71. Grichnik KP, Ferrante FM. The difference between acute and chronic pain. *Mt Sinai J Med* 1991;58:217-220.
72. Koh JS, Ko HJ, Wang SM, Cho KJ, Kim JC, Lee SJ, et al. Depression and somatic symptoms may influence on chronic prostatitis/chronic pelvic pain syndrome: a preliminary study. *Psychiatry Investig* 2014;11:495-499.
73. Bonezzi C, Demartini L, Buonocore M. Chronic pain: not only a matter of time. *Minerva Anestesiol* 2012;78:704-711.
74. Devon HA, Piano MR, Rosenfeld AG, Hoppensteadt DA. The association of pain with protein inflammatory biomarkers: a review of the literature. *Nurs Res* 2014;63:51-62.
75. Miller AH, Maletic V, Raison CL. Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biol Psychiatry* 2009;65:732-741.